## The Pelvic Pain and Urinary/Frequency (PUF)

**Symptom** 

Score

**Bother** 

Score

Patient Symptom Scale						
Please circle the answer that best describes how you feel for each question.						
	0	1	2	3	4	
1. How many times do you go to the bathroom during the day?	3–6	7–10	11–14	15–19	20+	
2a. How many times do you go to the bathroom at night?	0	1	2	3	4+	
2b. If you get up at night to go to the bathroom, does it bother you?	Never	Mildly	Moderate	Severe		
3. Are you currently sexually active? Yes No						

Never

Never

Never

Never

Never

Never

Occasionally

Occasionally

Occasionally

Occasionally

Mild

Occasionally

Mild

Occasionally

Usually

Usually

Usually

Usually

**Moderate** 

Usually

**Moderate** 

Usually

**Always** 

**Always** 

**Always** 

**Always** 

Severe

**Always** 

Severe

**Always** 

Symptom Score (1,  $2\alpha$ ,  $4\alpha$ , 5, 6,  $7\alpha$ ,  $8\alpha$ ) =

Total Score (Symptom Score + Bother Score) =

Bother Score (2b, 4b, 7b, 8b) =

4a. If you are sexually active, do you now or have you ever had pain or symptoms during or after sexual intercourse?

4b. If you have pain, does it make you avoid sexual intercourse?

Do you have urgency after going to the bathroom?

A total score of:

10-14= 74% likelihood of IC

15-19= 86% likelihood of IC 20 or above = 91% likelihood of IC

7a. If you have pain, is it usually...

8a. If you have urgency, is it usually...

8b. Does your urgency bother you?

7b. Does your pain bother you?

Do you have pain associated with your bladder or in your pelvis (vagina, lower abdomen, urethra, perineum, testes, or scrotum)?